





## **Further Reading**

The Royal Australian College of General Practitioners. Supporting smoking cessation: A guide for health professionals. 2nd edn. East Melbourne, Vic: RACGP, 2019.

Available to download: https://www.racgp.org.au/clinical-resources/clinical-guidelines/key-racgp-guidelines/view-all-racgp-guidelines/supporting-smoking-cessation

## References

- Coleman, T, Chamberlain, C, Davey, M, Cooper, SE, Leonardi-Bee, J, 2015, 'Pharmacological interventions for promoting smoking cessation during pregnancy', *Cochrane Database of Systematic Reviews*, no. 12.
- Dempsey, DA, & Benowitz, NL, 2001, 'Risks and Benefits of Nicotine to Aid Smoking Cessation in Pregnancy', *NL Drug-Safety*, vol.24, no.4, pp.277-322.
- Faber, M S and Fuhr, U, 2004, 'Time response of cytochrome P450 1A2 activity on cessation of heavy smoking', *Clinical Pharmacology & Therapeutics*, vol.76, no.2, pp.178–184.
- Ferguson, SG, Gitchell, JG, Shiffman, S, Sembower, MA, Rohay, JM, & Allen, J, 2011, 'Providing accurate safety information may increase a smoker's willingness to use nicotine replacement therapy as part of a quit attempt', *Addictive behaviors*, vol. 36, no. 7, pp. 713-16.
- Gould, G, Bittoun, R, & Clarke, M, 2015, 'A Pragmatic Guide for Smoking Cessation Counselling and the Initiation of Nicotine Replacement Therapy for Pregnant Aboriginal and Torres Strait Islander Smokers', *Journal of Smoking Cessation*, vol.10, no.2, pp.96-105.
- Greenhalgh, EM, Stillman, S, & Ford, C, 2016, 'Smoking and mental health', in MM. Scollo, and MH Winstanley, eds., *Tobacco in Australia: Facts and issues,* Melbourne: Cancer Council Victoria.
- Hartmann-Boyce, J, Stead, L F, Cahill, K, & Lancaster, T, 2014, 'Efficacy of interventions to combat tobacco addiction: Cochrane update of 2013 reviews', *Addiction*, vol.109, no.9, pp. 1414–1425
- Hartmann-Boyce, J, Stead, L F, Cahill, K, & Lancaster, T, 2013, 'Efficacy of interventions to combat tobacco addiction: Cochrane update of 2012 reviews', *Addiction*, vol.108, no.10, pp.1711–1721.
- Heatherton, T F Kozlowski, L T, Frecker, R C, Rickert, W, & Robinson, J, 1989, 'Measuring the Heaviness of Smoking: using self-reported time to the first cigarette of the day and number of cigarettes smoked per day', *British Journal of Addiction*, vol.84, no.7, pp.791–800.

## References cont.

- Hukkanen, J, Jacob, P, & Benowitz, NL, 2005, 'Metabolism and Disposition Kinetics of Nicotine', *Pharmacological Reviews*, vol.57, no.1, pp.79-115.
- Mendelsohn, CP, 2013, 'How to treat Nicotine dependence', *Australian Doctor*, www.australiandoctor.com.au
- Mendelsohn, CP, Kirby, DP, & Castle, DJ, 2015, 'Smoking and mental illness An update for psychiatrists', *Australasian Psychiatry*, vol.23, no.1, pp.37-43.
- Mendelsohn, CP, & Montabello, ME, 2013, 'Smokers with mental illness breaking down the myths, *Medicine Today*, vol.14, no.8, pp53-57.
- Mendelsohn CP, Gould, GS, Oncken, C, 2014, 'Management of smoking in pregnant women', Australian Family Physician, vol.43 no. 1/2, pp.46-51.
- Zwar, N, Richmond, R, Borland, R, Peters, M, Litt, J, Bell, J, Caldwell, B, Ferretter, I, 2011, Supporting smoking cessation: a guide for health professionals, Royal Australian College of General Practitioners, (updated July 2014).
- Shiffman, S, & Ferguson, S G, 2008, 'Nicotine patch therapy prior to quitting smoking: a meta-analysis', *Addiction*, vol.103, no.4, pp.557–563.
- Stead LF, Ferera, R, Bullen, C, Mant, D, Hartmann-Boyce, J, Cahill, K, Lancaster, T, 2012, 'Nicotine replacement therapy for smoking cessation (Review)', *Cochrane Database of Systematic Reviews*, no.11.
- Stead LF, Hartmann-Boyce, J, Ferera, R, Lancaster, T, 2013, 'Telephone counselling for smoking cessation (Review)', *Cochrane Database of Systematic Reviews*, no.8.
- Tyman, L, Bonevski, B, Paul, C, & Bryant, J, 2014, 'Perceived barriers to smoking cessation in selected vulnerable groups: a systematic review of the qualitative and quantitative literature', *BMJ Open*, vol.4, no.12.
- Williams, JM, Steinberg, ML, Griffiths, KG, & Cooperman, N, 2013, 'Smokers with behavioral health comorbidity should be designated a tobacco use disparity group', *American Journal of Public Health*, vol.103, no.9, pp.1549-1555.









